ELSEVIER

Toxicology 144 (2000) 243–245

TOXICOLOGY

www.elsevier.com/locate/toxicol

Contents of volume 144

Special Issue

FESTSCHRIFT DEDICATED TO PROFESSOR DR. K.J. NETTER

Guest Editor:
Edmund Maser

Laudatio

In honour of the past European editor: K.J. Netter
W. Legrum (Germany) 1–4

Articles

Studies on pyrazinoylguanidine: a novel antihypertensive, hypoglycemic and lipolytic drug intended for adjunctive use in hypertensive patients with type 2 diabetes mellitus
E.S. Vesell, K.H. Beyer, Jr. (USA) 5–11
Pyrazinolguanidine reduces elevated glucose, lipids and blood pressure in type 2 diabetes patients

Use of suppression-PCR subtractive hybridisation to identify genes that demonstrate altered expression in male rat and guinea pig livers following exposure to Wy-14,643, a peroxisome proliferator and non-genotoxic hepatocarcinogen
J.C. Rockett, K.E. Swales, D.J. Esdaile, G.G. Gibson (UK) 13–29
Differential gene display identifies genes up- and down- regulated by a peroxisome proliferator

Bioflavonoids: selective substrates and inhibitors for cytochrome P450 CYP1A and CYP1B1
H. Doostdar, M.D. Burke, R.T. Mayer (USA, UK) 31–38
Bioflavonoids: Selective substrates and inhibitors for cytochrome P450 CYP1A

Estimation of human exposure to styrene and ethylbenzene
W. Tang, I. Hemm, G. Eisenbrand (Germany) 39–50
Exposure of the consumer (non-smoking) to styrene and ethylbenzene results to >90% from inhaled air

Density-dependent growth of normal and nodular hepatocytes
K.W. Bock, H. Gschaidmeier, B.S. Bock-Hennig, L.C. Eriksson (Germany, Sweden) 51–56
Nodular hepatocytes do not show cell contact inhibition

Effect of selenium supplementation on the activities of glutathione metabolizing enzymes in human hepatoma Hep G2 cell line
M.H. Helmy, S.S. Ismail, H. Fayed, E.A. El-Bassiouni (Egypt) 57–61
Selenium may protect against oxidative stress in tissue culture

Heterogeneity of 11β-hydroxysteroid dehydrogenase type 1 microsomal carbonyl reductase (11β-HSD CR) in guinea pig tissues. Purification of the liver form suggests modification in the cosubstrate binding site
U.C.T. Oppermann, E. Möbus, G. Nagel, E. Maser (Sweden, Germany) 63–69
Isoforms of 11beta-HSD in guinea pig tissues

Molecular and structural aspects of xenobiotic carbonyl metabolizing enzymes. Role of reductases and dehydrogenases in xenobiotic phase I reactions
U.C.T. Oppermann, E. Maser (Sweden, Germany) 71–81
Involvement of dehydrogenases/reductases in phase I metabolism

Use of precision-cut tissue slices in organ culture to study metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) by hamster lung, liver and kidney
E. Richter, S. Friesenegger, J. Engl, A.R. Tricker (Germany, Switzerland) 83–91
No correlations between target organ carcinogenicity and metabolic activation of NNK in hamster tissues
Induction of cytochrome P450 2B1-mRNA and pentoxyresorufin O-depentylation after exposure of precision-cut rat liver slices to phenobarbital
D. Müller, P. Steinmetzer, K. Pissowotzki, R. Glöckner (Germany) 93–97
In vitro exposure of liver slices to phenobarbital induces CYP2B1-mRNA tremendously
Effects of cotinine at cholinergic nicotinic receptors of the sympathetic superior cervical ganglion of the mouse
K.-C. Schroff, P. Lovich, O. Schmitz, S. Aschhoff, E. Richter, J. Remien (Germany) 99–105
Cotinine has antagonistic action at peripheral neuronal nicotinic receptors
Environmental inhibition of 11β-hydroxysteroid dehydrogenase
M.M. Reidenberg (USA) 107–111
Dietary and other exogeneous inhibitors of 11βHSD may influence electrolyte metabolism
Human 11β-hydroxysteroid dehydrogenase 1/carbonyl reductase: recombinant expression in the yeast *Pichia pastoris* and *Escherichia coli*
A. Blum, H.-J. Martin, E. Maser (Germany) 113–120
Recombinant 11ß-HSD1 serves as a valuable tool in studies on carbonyl compound detoxification
Toxicologic evidence of developmental neurotoxicity of environmental chemicals
H.R. Andersen, J.B. Nielsen, P. Grandjean (Denmark) 121–127
Basic kinetic data are lacking in risk assessment of developmental neurotoxicants
Genetic polymorphism of CYP2A6 in the German population
M. Bourian, H. Gullstén, W. Legrum (Germany, Finland) 129–137
Genetic polymorphism of CYP2A6
CYP2A6: a human coumarin 7-hydroxylase
O. Pelkonen, A. Rautio, H. Raunio, M. Pasanen (Finland) 139–147
Substrates, inhibitors and modifying factors of human CYP2A6 are described
Laryngeal mucosa of head and neck cancer patients shows increased DNA damage as detected by single cell microgel electrophoresis
P. Schmezer, T. Rupprecht, M. Tisch, H. Maier, H. Bartsch (Germany) 149–154
Head and neck cancer patients show increased DNA damage in their laryngal mucosa
Confounding variables in the environmental toxicology of arsenic
T. Gebel (Germany) 155–162
Se, Sb, Zn, cellular resistance and genetic polymorphisms may modulate the long-term toxicity of As
Lack of quinone reductase activity suggests that amyloid-β peptide/ERAB induced lipid peroxidation is not directly related to production of reactive oxygen species by redoxcycling
S. Salim, C. Filling, E. Mårtensson, U.C.T. Oppermann (Sweden) 163–168
Oxidative stress by amyloid beta peptide/enzyme interactions
Dose and time as variables of toxicity
K.K. Rozman, J. Doull (USA, Germany) 169–178
Development of a theory of toxicology
Tissue distribution of 2-(2-nitro-4-trifluoromethylbenzoyl)-cyclohexane-1,3-dione (NTBC) and its effect on enzymes involved in tyrosine catabolism in the mouse
E.A. Lock, P. Gaskin, M.K. Ellis, W. McLean Provan, M. Robinson, L.L. Smith (UK) 179–187
Inhibition of mouse liver HPPD causes tyrosinaemia, but not corneal injury
Modulation of daunorubicin toxicity by liposomal encapsulation and use of specific inhibitors in vitro
T. Iffert, M. Soldan, A. Moeller, E. Maser (Germany) 189–195
Liposomal encapsulation did not increase DRC toxicity
Structural characteristics of human P450s involved in drug metabolism: QSARs and lipophilicity profiles
D.F.V. Lewis (UK) 197–203
QSARs in human cytochromes P450
Reduction of gastrointestinal toxicity of NSAIDs via molecular modifications leading to antioxidant anti-inflammatory drugs
P.N. Kourounakis, K. Tsiakitzis, A.P. Kourounakis, D. Galanakis (Greece) 205–210
Antioxidant NSAID derivatives with greatly reduced ulcerogenicity
Enhanced expression of CYP1B1 in *Escherichia coli*
I. Jansson, I. Stoilov, M. Sarfarazi, J.B. Schenkman (USA) 211–219
Human cytochrome P450 expression in Escherichia coli

Reversal of HER-2 over-expression renders human ovarian cancer cells highly resistant to taxol
A. Aigner, S.S. Hsieh, C. Malerczyk, F. Czubayko (Germany, USA) 221–228
Taxol cytotoxicity related to HER-2 expression level
Monitoring of cytochrome P-450 1A activity by determination of the urinary pattern of caffeine metabolites in Wistar and hyperbilirubinemic Gunn rats
U. Jorritsma, E. Schrader, G. Klaunick, J. Kapitulnik, K.I. Hirsch-Ernst, G.F. Kahl, H. Foth (Germany, Israel) 229–236
The caffeine metabolite pattern was used to assess CYP1A induction

Author index of volume 144 237
Subject index of volume 144 239–241
Contents of volume 144 243–245